

Reductive whole-cell biotransformation with *Corynebacterium glutamicum*: improvement of NADPH generation from glucose by a cyclized pentose phosphate pathway using *pfkA* and *gapA* deletion mutants

Solvej Siedler · Steffen N. Lindner · Stephanie Bringer · Volker F. Wendisch · Michael Bott

Received: 24 May 2012 / Revised: 16 July 2012 / Accepted: 16 July 2012 / Published online: 1 August 2012

© The Author(s) 2012. This article is published with open access at Springerlink.com

Abstract In this study, the potential of *Corynebacterium glutamicum* for reductive whole-cell biotransformation is shown. The NADPH-dependent reduction of the prochiral methyl acetoacetate (MAA) to the chiral (*R*)-methyl 3-hydroxybutyrate (MHB) by an alcohol dehydrogenase from *Lactobacillus brevis* (*Lbadh*) was used as model reaction and glucose served as substrate for the regeneration of NADPH. Since NADPH is mainly formed in the oxidative branch of the pentose phosphate pathway (PPP), *C. glutamicum* was engineered to redirect carbon flux towards the PPP. Mutants lacking the genes for 6-phosphofructokinase (*pfkA*) or glyceraldehyde 3-phosphate dehydrogenase (*gapA*) were constructed and analyzed with respect to growth, enzyme activities, and biotransformation performance. Both mutants showed strong growth defects in glucose minimal medium. For biotransformation of MAA to MHB using glucose as reductant, strains were transformed with an *Lbadh* expression plasmid. The wild type showed a specific MHB production rate of $3.1 \text{ mmol}_{\text{MHB}} \text{ h}^{-1} \text{ g}_{\text{cdw}}^{-1}$ and a yield of $2.7 \text{ mol}_{\text{MHB}} \text{ mol}_{\text{glucose}}^{-1}$. The ΔpfkA mutant showed a similar MHB production rate, but reached a yield of $4.8 \text{ mol}_{\text{MHB}} \text{ mol}_{\text{glucose}}^{-1}$, approaching the maximal value of $6 \text{ mol}_{\text{NADPH}} \text{ mol}_{\text{glucose}}^{-1}$ expected for a partially cyclized PPP. The specific biotransformation rate of the ΔgapA mutant was decreased by 62 %

compared to the other strains, but the yield was increased to $7.9 \text{ mol}_{\text{MHB}} \text{ mol}_{\text{glucose}}^{-1}$, which to our knowledge is the highest one reported so far for this mode of NADPH regeneration. As one fourth of the glucose was converted to glycerol, the experimental yield was close to the theoretically maximal yield of $9 \text{ mol}_{\text{NADPH}} \text{ mol}_{\text{glucose}}^{-1}$.

Keywords *Corynebacterium glutamicum* · Pathway engineering · NADPH yield · Pentose phosphate pathway · Resting cells · Reductive whole-cell biotransformation · Phosphofructokinase · Glyceraldehyde 3-phosphate dehydrogenase · *pfk* · *gap*

Introduction

Whole-cell biotransformation has become an important method in chemoenzymatic synthesis, e.g., for the production of amino acids and chiral alcohols (Ishige et al. 2005). *Corynebacterium glutamicum* is a Gram-positive, non-pathogenic soil bacterium which is predominantly used for the large-scale industrial production of the flavor enhancer L-glutamate and the food additive L-lysine (Pfefferle et al. 2003; Kimura 2003; Hermann 2003). Recent metabolic engineering studies have shown that *C. glutamicum* is also capable of producing a variety of other commercially interesting compounds, e.g., other L-amino acids (Wendisch et al. 2006), D-amino acids (Stäbler et al. 2011), organic acids such as succinate (Okino et al. 2008; Litsanov et al. 2012a, b), diamines such as cadaverine (Mimitsuka et al. 2007) or putrescine (Schneider and Wendisch 2010), biofuels such as ethanol or isobutanol (Inui et al. 2004; Smith et al. 2010; Blombach et al. 2011), or proteins (Meissner et al. 2007). An overview of the product spectrum of *C. glutamicum* can be found in a recent review (Becker and Wittmann 2011).

Solvej Siedler and Steffen N. Lindner contributed equally to this work.

S. Siedler · S. Bringer (✉) · M. Bott (✉)
Institut für Bio- und Geowissenschaften, IBG-1: Biotechnologie,
Forschungszentrum Jülich GmbH,
52425 Jülich, Germany
e-mail: st.bringer-meyer@fz-juelich.de
e-mail: m.bott@fz-juelich.de

S. N. Lindner · V. F. Wendisch
Faculty of Biology & CeBiTec, Bielefeld University,
33615 Bielefeld, Germany

In this study, we analyzed *C. glutamicum* mutants lacking either *pfkA* or *gapA* for their behavior in reductive whole-cell biotransformation. The results supported the view that the PPP operates in cyclic manner, oxidizing glucose to CO₂ with concomitant reduction of NADP⁺ to NADPH.

Materials and methods

Chemicals and enzymes

Chemicals were obtained from Sigma-Aldrich (Taufkirchen, Germany), Qiagen (Hilden, Germany), Merck (Darmstadt, Germany), and Roche Diagnostics (Mannheim, Germany).

Bacterial strains, plasmids, media, and growth conditions

Strains and plasmids used in this work are listed in Table 1. *E. coli* strains were transformed by the method described by Hanahan (1983) and cultivated in LB medium (Miller 1972). *E. coli* DH5 α was used for cloning purposes and *C. glutamicum* ATCC 13032 and derivatives for gene expression and whole-cell biotransformation. When required, antibiotics were added to the medium at a final concentration of 50 μ g kanamycin ml⁻¹ (pEKEx2-LbADH) or 100 μ g spectinomycin ml⁻¹ (pEKEx3 derivatives).

For growth experiments with *C. glutamicum*, 50-ml LB overnight cultures were inoculated from LB plates, harvested by centrifugation (10 min, 3,220 \times g), washed in CgXII medium (Eggeling and Bott 2005), and inoculated in CgXII medium containing 100 mM glucose to a final optical density at 600 nm (OD₆₀₀) of 1. When appropriate, 1 mM isopropyl- β -D-thiogalactopyranosid (IPTG), 25 μ g ml⁻¹ kanamycin, and 100 μ g ml⁻¹ spectinomycin was added. For all growth experiments, 500 ml baffled shake flasks with 50 ml CgXII medium were used and incubated at 30 °C and 120 rpm. Growth was followed by OD₆₀₀ determination using a UV-1650 PC photometer (Shimadzu, Duisburg, Germany). The biomass concentration was calculated from OD₆₀₀ values using an experimentally determined correlation factor of 0.25 g (dry weight) of cells (cdw) per liter for an OD₆₀₀ of 1 (Kabus et al. 2007). For the determination of enzyme activity in cell-free extracts, 50 ml LB medium containing 1 mM IPTG and 100 μ g ml⁻¹ spectinomycin was inoculated from LB overnight cultures to an OD₆₀₀ of 0.5. At an OD₆₀₀ of 4, cells were harvested by centrifugation (10 min, 3,220 \times g, 4 °C) and stored at -20 °C until use.

Recombinant DNA work

Standard methods like polymerase chain reaction (PCR), restriction, or ligation were carried out according to established

protocols (Sambrook and Russell 2001). *E. coli* cells were transformed by the CaCl₂ method (Hanahan et al. 1991). DNA sequencing was performed by Eurofins MWG Operon (Germany). Oligonucleotides (listed in Table 2) were synthesized by Biolegio bv (Nijmegen, The Netherlands) and Eurofins MWG Operon (Germany).

Construction of deletion mutants and plasmids

C. glutamicum deletion mutants were constructed using pK19*mobsacB* (Schäfer et al. 1994) using the procedure described by Niebisch and Bott (2001). Upstream and downstream flanking regions of *pfkA* (cg1409), and *gapA* (cg1791) were amplified by PCR using the oligonucleotide pairs *pfkA*-Del-A/*pfkA*-Del-B and *pfkA*-Del-C/*pfkA*-Del-D for deletion of *pfkA*, and *gapA*-Del-A/*gapA*-Del-B and *gapA*-Del-C/*gapA*-Del-D for deletion of *gapA* (see Table 2 for primer sequences). The upstream and downstream flanking regions of each gene were fused by overlap extension PCR, resulting in a DNA fragment of about 1 kb. The resulting PCR products were cloned into SmaI-restricted vector pK19*mobsacB* resulting in pK19*mobsacB* Δ *pfkA*, and pK19*mobsacB* Δ *gapA*. The correctness of the cloned PCR fragments was confirmed by DNA sequencing. Transformation of *C. glutamicum* wild type with these plasmids and selection for the first and second homologous recombination was performed as described (Niebisch and Bott 2001; Rittmann et al. 2003). Kanamycin-sensitive and sucrose-resistant clones were analyzed by PCR using oligonucleotide pairs *pfkA*-Del-Ver-fw/*pfkA*-Del-Ver-rv or *gapA*-Del-Ver-fw/*gapA*-Del-Ver-rv.

For the complementation of deletion mutants, the genes *pfkA* (cg1409), and *gapA* (cg1791) from *C. glutamicum* and the genes *pfkA* (b3916) and *pfkB* (b3916) from *E. coli* were amplified via PCR from genomic DNA of *C. glutamicum* WT, which was prepared as described previously (Eikmanns et al. 1995), and *E. coli* MG1655 genomic DNA, which was prepared by using the DNA isolation kit (Roche, Mannheim, Germany). PCR was performed using the following oligonucleotide pairs: *pfkA*-cgl-fw/*pfkA*-cgl-rv, *gapA*-cgl-fw/*gapA*-cgl-rv, *pfkA*-eco-fw/*pfkA*-eco-rv, and *pfkB*-eco-fw/*pfkB*-eco-rv (see Table 2). To allow IPTG-inducible expression of *pfkA*, and *gapA* from *C. glutamicum* and *pfkA*, and *pfkB* from *E. coli* the corresponding PCR products were ligated into the SmaI-restricted vector pEKEx3 resulting in pEKEx3-*pfkA*^{Cgl}, pEKEx3-*gapA*^{Cgl}, pEKEx3-*pfkA*^{Eco}, and pEKEx3-*pfkB*^{Eco}.

For the construction of the expression plasmid pEKEx2-*Lbadh*, the *adh* gene of *L. brevis* was amplified together with a 9-bp linker and an artificial ribosome binding site (AAGGAG) using the oligonucleotides *Lbadh*_for and *Lbadh*_rev and the plasmid pBtac*Lbadh* as template (Ernst et al. 2005). The PCR product was digested with *Bam*HI and *Eco*RI and cloned into the vector pEKEx2. The correctness

Table 1 Strains and plasmids used in this work

Strains and plasmids	Relevant characteristics	Reference
Strains		
<i>E. coli</i> DH5 α	F [−] ϕ 80 Δ lacZ Δ M15 Δ (lacZYA-argF) U169 <i>deoR recA1 endA1 hsdR17</i> (rk [−] , mk ⁺) <i>phoA supE44</i> λ^{-} <i>thi-1 gyrA96 relA1</i>	(Hanahan 1983), Invitrogen
<i>C. glutamicum</i> ATCC13032	Wild type, biotin auxotrophic	(Abe et al. 1967)
Δ <i>pfkA</i>	<i>C. glutamicum</i> ATCC13032 Δ <i>pfkA</i> (cg1409)	This study
Δ <i>gapA</i>	<i>C. glutamicum</i> ATCC13032 Δ <i>gapA</i> (cg1791)	This study
WT/pEKEx3	<i>C. glutamicum</i> ATCC13032 with pEKEx3	This study
WT/pEKEx3- <i>pfkA</i> ^{Cgl}	<i>C. glutamicum</i> ATCC13032 with pEKEx3- <i>pfkA</i> ^{Cgl}	This study
WT/pEKEx3- <i>pfkA</i> ^{Eco}	<i>C. glutamicum</i> ATCC13032 with pEKEx3- <i>pfkA</i> ^{Eco}	This study
WT/pEKEx3- <i>pfkB</i> ^{Eco}	<i>C. glutamicum</i> ATCC13032 with pEKEx3- <i>pfkB</i> ^{Eco}	This study
WT/pEKEx3- <i>gapA</i> ^{Cgl}	<i>C. glutamicum</i> ATCC13032 with pEKEx3- <i>gapA</i> ^{Cgl}	This study
WT/pEKEx2- <i>Lbadh</i>	<i>C. glutamicum</i> ATCC13032 with pEKEx2- <i>Lbadh</i>	This study
Δ <i>pfkA</i> /pEKEx3	<i>C. glutamicum</i> ATCC13032 Δ <i>pfkA</i> with pEKEx3	This study
Δ <i>pfkA</i> /pEKEx3- <i>pfkA</i> ^{Cgl}	<i>C. glutamicum</i> ATCC13032 Δ <i>pfkA</i> with pEKEx3- <i>pfkA</i> ^{Cgl}	This study
Δ <i>pfkA</i> /pEKEx3- <i>pfkA</i> ^{Eco}	<i>C. glutamicum</i> ATCC13032 Δ <i>pfkA</i> with pEKEx3- <i>pfkA</i> ^{Eco}	This study
Δ <i>pfkA</i> /pEKEx3- <i>pfkB</i> ^{Eco}	<i>C. glutamicum</i> ATCC13032 Δ <i>pfkA</i> with pEKEx3- <i>pfkB</i> ^{Eco}	This study
Δ <i>pfkA</i> /pEKEx2- <i>Lbadh</i>	<i>C. glutamicum</i> ATCC13032 Δ <i>pfkA</i> with pEKEx2- <i>Lbadh</i>	This study
Δ <i>gapA</i> /pEKEx3	<i>C. glutamicum</i> ATCC13032 Δ <i>gapA</i> with pEKEx3	This study
Δ <i>gapA</i> /pEKEx3- <i>gapA</i> ^{Cgl}	<i>C. glutamicum</i> ATCC13032 Δ <i>gapA</i> with pEKEx3- <i>gapA</i> ^{Cgl}	This study
Δ <i>gapA</i> /pEKEx2- <i>Lbadh</i>	<i>C. glutamicum</i> ATCC13032 Δ <i>gapA</i> with pEKEx2- <i>Lbadh</i>	This study
Plasmids		
pEKEx2	Kan ^r ; <i>E. coli</i> – <i>C. glutamicum</i> shuttle vector for regulated gene expression (P _{tac} <i>lacI</i> ^q pBL1 <i>oriV</i> _{C.g.} pUC18 <i>oriV</i> _{E.c.})	(Eikmanns et al. 1991)
pEKEx2- <i>Lbadh</i>	Kan ^r ; pEKEx2 derivative with <i>adh</i> gene from <i>Lactobacillus brevis</i>	This study
pEKEx3	Spec ^r ; <i>C. glutamicum</i> / <i>E. coli</i> shuttle vector (P _{tac} , <i>lacI</i> ^q ; pBL1, <i>oriV</i> _{C.g.} , <i>oriV</i> _{E.c.})	(Stansen et al. 2005)
pEKEx3- <i>pfkA</i> ^{Cgl}	Spec ^r ; derivative of pEKEx3 for regulated expression of <i>pfkA</i> (cg1409) of <i>C. glutamicum</i>	This study
pEKEx3- <i>gapA</i> ^{Cgl}	Spec ^r ; derivative of pEKEx3 for regulated expression of <i>gapA</i> (cg1791) of <i>C. glutamicum</i>	This study
pEKEx3- <i>pfkA</i> ^{Eco}	Spec ^r ; derivative of pEKEx3 for regulated expression of <i>pfkA</i> (b3916) of <i>E. coli</i>	This study
pEKEx3- <i>pfkB</i> ^{Eco}	Spec ^r ; derivative of pEKEx3 for regulated expression of <i>pfkB</i> (b1723) of <i>E. coli</i>	This study
pK19 <i>mobsacB</i>	Kan ^r ; mobilizable <i>E. coli</i> vector used for the construction of <i>C. glutamicum</i> insertion and deletion mutants (RP4 <i>mob</i> ; <i>sacB</i> _{B.sub.} ; <i>lacZ</i> α ; <i>oriV</i> _{E.c.})	(Schäfer et al. 1994)
pK19 <i>mobsacB</i> Δ <i>pfkA</i>	Kan ^r ; pK19 <i>mobsacB</i> derivative containing a PCR product which covers the flanking regions of the <i>C. glutamicum</i> <i>pfkA</i> (cg1409) gene	This study
pK19 <i>mobsacB</i> Δ <i>gapA</i>	Kan ^r ; pK19 <i>mobsacB</i> derivative containing a PCR product which covers the flanking regions of the <i>C. glutamicum</i> <i>gapA</i> (cg1791) gene	This study

of the cloned PCR fragments in the plasmids was confirmed by DNA sequencing.

Enzyme activity assays

For the determination of alcohol dehydrogenase activity, cells were harvested by centrifugation (10,000 \times g, 4 °C, 5 min) 30 min after start of biotransformation and stored at −20 °C until use. The cells were resuspended in 100 mM potassium phosphate buffer, pH 6.5, with 1 mM dithiothreitol and 1 mM MgCl₂. Cells were disrupted at 4 °C by 3 \times 15 s bead-beating with 0.1-mm-diameter glass beads using a Silamat S5 (Ivoclar Vivadent GmbH, Germany) and crude extracts were centrifuged at 16,000 \times g (4 °C, 20 min) to

remove intact cells and cell debris. The supernatants were used as cell-free extracts. Alcohol dehydrogenase activity was determined photometrically at 340 nm using a mixture of 10 mM methyl acetoacetate, 250 μ M NADPH, and 1 mM MgCl₂ in 100 mM potassium phosphate buffer, pH 6.5. The reactions were started by adding different dilutions of the cell-free extract. For rate calculation, an extinction coefficient for NADPH at 340 nm of 6.22 mM^{−1} cm^{−1} was used. One unit of enzyme activity corresponds to 1 μ mol NADPH consumed per minute.

For the determination of the specific activity of phosphofructokinase and glyceraldehyde 3-phosphate dehydrogenase, cells were harvested by centrifugation (3,220 \times g, 4 °C, 10 min) and washed in the appropriate buffer (see below)

Table 2 Sequences of oligonucleotide primers

Name	Sequence (5'–3')	Function and relevant characteristics
pfkA-cgl-fw	GGATCCGAAAGGAGGCCCTTCAGATGGAAGACATGCGAATTGCTAC	OE of Cgl <i>pfkA</i> ; start; BamHI ; RBS
pfkA-cgl-rv	GGATCCCTATCCAAACATTGCCTGGGC	OE of Cgl <i>pfkA</i> ; stop; BamHI
gapA-cgl-fw	<i>AAGGAGATATAGATATGACCATTTCGTGTTGGTATTAAC</i>	OE of Cgl <i>gapA</i> ; start; RBS
gapA-cgl-rv	<i>TTAGAGCTTGGGAAGCTACGAGCTC</i>	OE of Cgl <i>gapA</i> ; stop
pfkA-eco-fw	CCGGATCCGAAAGGAGGCCCTTCAGATGATTAAGAAAATCGGTGTGTTGAC	OE of Eco <i>pfkA</i> ; start; BamHI ; RBS
pfkA-eco-rv	CCGGATCCCTTAATACAGTTTTTTCGCGCAGTC	OE of Eco <i>pfkA</i> ; stop; BamHI
pfkB-eco-fw	GACTGCAGGAAAGGAGGCCCTTCAGATGGTACGTATCTATACGTTGACAC	OE of Eco <i>pfkB</i> ; start; PstI ; RBS
pfkB-eco-rv	GGCTGCAGTTAGCGGGAAAGGTAAGCGTAA	OE of Eco <i>pfkB</i> ; stop; PstI
pfkA-Del-A	CCGGAATATCTCGACGCCACAGAACGC	Del of <i>pfkA</i>
pfkA-Del-B	<i>CCCATCCACTAAACTTAAACAATTCGCATGTCTTCCATATTAAACCCATCACAAACACCCGC</i>	Del of <i>pfkA</i> ; linker sequence
pfkA-Del-C	<i>TGTTTAAGTTTAGTGGATGGGGAACGCTGGGTACTGCCAGGCAATGTTT</i>	Del of <i>pfkA</i> ; linker sequence
pfkA-Del-D	CCGAAGGAATAGACGAGTTAACAAAACCTACGGTCTG	Del of <i>pfkA</i>
pfkA-Del-Ver-fw	GCCAAACTCGAGTAGCCCGG	Verification of <i>pfkA</i> Del
pfkA-Del-Ver-rv	CCACAGCTTCAGTCATGCCC	Verification of <i>pfkA</i> Del
gapA-Del-A	GGCTGATCCTCAAATGACCAAG	Del of <i>gapA</i>
gapA-Del-B	<i>CCCATCCACTAAACTTAAACAACCAACACGAATGGTCATGTTG</i>	Del of <i>gapA</i> ; linker sequence
gapA-Del-C	<i>TGTTTAAGTTTAGTGGATGGGCTGCGTCTGACCGAGCTCGTAG</i>	Del of <i>gapA</i> ; linker sequence
gapA-Del-D	CACCGAAGCCGTCAGAAACGAATG	Del of <i>gapA</i>
gapA-Del-Ver-fw	CCAACTTCGACGATGCCAATC	Verification of <i>gapA</i> Del
gapA-Del-Ver-rv	CTCTGGTGATTCTGCGATCTTTTC	Verification of <i>gapA</i> Del
lbADH_for	CAGTGGATCCGAAAGGAGGCCCTTCAGATGCTAACCGTTTGGATGG	OE of <i>Lb adh</i> ; start; BamHI ; RBS
lbADH_rev	GTCTGAATTCTATTGAGCAGTGTAGCCACC	OE of <i>Lb adh</i> ; stop; EcoRI

Restriction sites are highlighted in bold; linker sequences for crossover PCR and ribosomal binding sites are shown in italics; stop and start codons are underlined

OE overexpression, Del deletion, RBS ribosomal binding site, Cgl *C. glutamicum*, Eco *E. coli*

and stored at -20°C until use. Cells were resuspended in 1 ml of the buffer and cell-free extracts were prepared by sonification as described previously (Stansen et al. 2005). All enzyme activity measurements were carried out at 30°C . Protein concentrations were determined with bovine serum albumin as standard using Bradford reagents (Sigma, Taufkirchen, Germany).

6-Phosphofructokinase activity was measured spectrophotometrically at 340 nm according to Babul (1978) by a coupled enzymatic assay with pyruvate kinase and lactate dehydrogenase. ADP formed in kinase reaction was used to convert phosphoenolpyruvate to pyruvate, which was subsequently

reduced to lactate with concomitant oxidation of NADH to NAD^{+} . The assay solution contained 100 mM Tris-HCl pH 7.5, 0.2 mM NADH, 1 mM ATP, 10 mM MgCl_2 , and 0.2 mM phosphoenolpyruvate. One unit of enzyme activity corresponds to 1 μmol NADH oxidized per minute.

Glyceraldehyde 3-phosphate dehydrogenase activity was measured according to Omumasaba et al. (2004). The assay contained 1 mM NAD^{+} , 50 mM Na_2HPO_4 , 0.2 mM EDTA, and 0.5 mM glyceraldehyde 3-phosphate in 50 mM triethanolamine hydrochloride (TEA) buffer pH 8.5. One unit of enzyme activity corresponds to 1 μmol NADH formed per minute.

Whole-cell biotransformation

For cultivation of the different recombinant *C. glutamicum* strains carrying the pEKEx2-*Lbadh* plasmid, a single colony of each strain was inoculated into 10 ml BHIS medium (37 g l⁻¹ brain heart infusion, 91 g l⁻¹ sorbitol) containing the appropriate selection marker as described above and grown overnight at 30 °C and 120 rpm. These pre-cultures were used for inoculation of the main cultures to an optical density at 600 nm (OD₆₀₀) of 0.4. Main cultures were grown in 100 ml BHIS medium in shake flasks in the presence of the appropriate selection marker and 0.5 mM IPTG at 30 °C and 120 rpm. The cells were harvested at an OD₆₀₀ between 2.5 and 5 by centrifugation (4,000×g, 4 °C, 7 min) and resuspended in a solution containing 111 mM glucose, 2 mM MgSO₄, and 250 mM potassium phosphate buffer, pH 6.5, to a cell density of 3 g_{cdw} l⁻¹. The biotransformation was started by adding 50 mM MAA and conducted in shake flasks at 30 °C and 120 rpm to prevent cell sedimentation. Specific productivities (mmol_{MHB} h⁻¹ g_{cdw}⁻¹) were determined by taking samples at 30–60-min time intervals over a period of 3 h. MHB and glucose concentrations of the samples were determined (see below). Specific productivities were calculated by dividing the slope of graphs showing MHB concentration vs. time by the cell dry weight, which remained constant.

Analysis of substrates and products

Methyl acetoacetate (MAA), (*R*)-methyl 3-hydroxybutyrate (MHB), glucose, and extracellular metabolites were analyzed by HPLC as described previously (Siedler et al. 2011).

Results

Growth behavior and in vitro enzyme activities of *C. glutamicum* wild-type and mutant strains

In a *C. glutamicum* mutant lacking 6-phosphofructokinase, glucose catabolism is forced to proceed via the pentose phosphate pathway. Fructose 6-phosphate formed in the PPP by transaldolase or transketolase has to re-enter the oxidative part of the PPP again and only glyceraldehyde 3-phosphate can be catabolized further via the lower part of the glycolytic pathway. Thus, the initial part of glucose catabolism in a $\Delta pfkA$ mutant can be described by the following equation: Glucose 6-phosphate + 6 NADP⁺ → Glyceraldehyde 3-phosphate + 3 CO₂ + 6 NADPH + 6 H⁺. Thus, 6 mol NADPH are formed per mole of glucose.

The deletion of the *pfkA* gene prevented growth in CgXII medium with 100 mM glucose (Table 3). The growth defect of the $\Delta pfkA$ mutant was complemented to levels of the WT

control (0.32 h⁻¹) by plasmid-based overexpression of either the homologous *pfkA* gene from *C. glutamicum* (0.32 h⁻¹) or of the heterologous *pfkA* gene from *E. coli* (0.33 h⁻¹) and increased to 0.16 h⁻¹ by heterologous expression of *pfkB* from *E. coli*. The slow growth of $\Delta pfkA$ /pEKEx3-*pfkB*^{Eco} was accompanied by a significantly higher biomass yield of 10.8 g l⁻¹ compared to 8.4 g l⁻¹ of WT/pEKEx3 or 8.6 g l⁻¹ of strain $\Delta pfkA$ /pEKEx3-*pfkA*^{Cgl}.

6-Phosphofructokinase activity was absent in the *pfkA* deletion strain (Table 3). Plasmid-borne expression of *C. glutamicum pfkA* or of *E. coli pfkA* or *pfkB* increased phosphofructokinase activity in the WT background from 0.04 U mg⁻¹ to 0.12, 0.11, and 0.19 U mg⁻¹, respectively. In the $\Delta pfkA$ background, phosphofructokinase activities of 0.10 to 0.13 U mg⁻¹ were determined when either *C. glutamicum pfkA* or *E. coli pfkA* or *pfkB* was overexpressed (Table 3).

C. glutamicum possesses two glyceraldehyde 3-phosphate dehydrogenases, GapA and GapB, but only GapA functions in the glycolytic direction as a $\Delta gapA$ deletion mutant was unable to grow in glucose minimal medium whereas a $\Delta gapB$ mutant showed no growth defect under these conditions (Omumasaba et al. 2004). A complete block of glyceraldehyde 3-phosphate conversion to 1,3-bisphosphoglycerate should lead to a complete oxidation of glucose in the PPP according to the equation: Glucose + 6 H₂O + 12 NADP⁺ → 6 CO₂ + 12 NADPH + 12 H⁺.

In agreement with previous results (Omumasaba et al. 2004), a deletion of the *gapA* gene in strain ATCC 13032 resulted in an inability to grow in glucose minimal medium. This defect was complemented by plasmid-based overexpression of the *gapA* gene. NAD⁺-dependent glyceraldehyde-3-phosphate dehydrogenase activity of cell-free extracts was 0.15 U mg⁻¹ in WT/pEKEx3

Table 3 Growth rates (μ) and biomass concentrations [cell dry weight (cdw) g l⁻¹] in glucose minimal medium with 1 mM IPTG and 100 μ g ml⁻¹ spectinomycin, and specific phosphofructokinase (Pfk) activity in cell extracts of the indicated *C. glutamicum* strains after cultivation in LB medium with 1 mM IPTG and 100 μ g ml⁻¹ spectinomycin

<i>C. glutamicum</i>	μ (h ⁻¹)	cdw (g l ⁻¹) ^a	Pfk activity (μmol min ⁻¹ mg ⁻¹)
WT/pEKEx3	0.32±0.00	8.43±0.18	0.04±0.01
WT/pEKEx3- <i>pfkA</i> ^{Cgl}	0.30±0.00	8.13±0.07	0.12±0.02
WT/pEKEx3- <i>pfkA</i> ^{Eco}	0.32±0.00	7.53±0.02	0.11±0.02
WT/pEKEx3- <i>pfkB</i> ^{Eco}	0.32±0.00	8.48±0.03	0.19±0.02
$\Delta pfkA$ /pEKEx3	0.00±0.00	0.14±0.01 ^b	0.00±0.00
$\Delta pfkA$ /pEKEx3- <i>pfkA</i> ^{Cgl}	0.32±0.01	8.63±0.07	0.10±0.01
$\Delta pfkA$ /pEKEx3- <i>pfkA</i> ^{Eco}	0.33±0.00	7.93±0.33	0.10±0.02
$\Delta pfkA$ /pEKEx3- <i>pfkB</i> ^{Eco}	0.16±0.00	10.80±0.10	0.13±0.01

^a Determination of cdw at maximal biomass

^b Determination of cdw after 24 h

and absent in strain $\Delta gapA/pEKEx3$. In strains WT/ $pEKEx3-gapA$ and $\Delta gapA/pEKEx3-gapA$, the glyceraldehyde 3-phosphate dehydrogenase activity with NAD^+ was found to be 0.26 and 0.13 $U\ mg^{-1}$, respectively (Table 4).

Biotransformation of MAA to MHB with the reference and the mutant strains

For biotransformation of MAA to MHB, the gene encoding the (R)-specific alcohol dehydrogenase of *L. brevis* (*Lbadh*) was overexpressed in *C. glutamicum* WT and in the deletion strains $\Delta pfkA$ and $\Delta gapA$ using plasmid $pEKEx2-Lbadh$. The specific NADPH-dependent MAA dehydrogenase activity in cell-free extracts of these strains was similar, ranging from 0.51 to 0.76 $U\ mg^{-1}$ in independent experiments. Assuming that the in vivo activities are comparable, they are not limiting the biotransformation rate. The *C. glutamicum* wild type showed a MAA dehydrogenase activity below 0.01 $U\ mg^{-1}$ with either NADPH or NADH as cofactor indicating that the biotransformation occurred only in the presence of the recombinant ADH from *L. brevis*. For the biotransformation, the strains were cultivated in BHIS medium to the exponential growth phase and then harvested and resuspended in 250 mM potassium phosphate buffer pH 6.5 containing 111 mM glucose and 2 mM $MgSO_4$ to a cell density of 3 $g_{cdw}\ l^{-1}$. The resulting cell suspensions were incubated at 30 °C and 120 rpm and the biotransformation was started by adding 50 mM MAA.

The kinetics of MHB production and of glucose consumption of the wild-type and the two mutant strains carrying $pEKEx2-Lbadh$ over a period of 180 min are shown in Fig. 2, and the rates and yields are listed in Table 5. It is evident from Fig. 2 that the rates of MHB production and glucose consumption were almost constant within the time period investigated and proportional to each other. The strain WT/ $pEKEx2-Lbadh$ showed an MHB production rate

of 3.14 $mmol\ h^{-1}\ g_{cdw}^{-1}$ and a glucose consumption rate of 1.17 $mmol\ h^{-1}\ g_{cdw}^{-1}$. This resulted in a MHB yield of 2.7 mol per mole of glucose, corresponding to an NADPH yield of 2.7 mol per mole of glucose. The strain $\Delta pfkA/pEKEx2-Lbadh$ had an 8 % reduced MHB production rate and a 49 % reduced glucose consumption rate, resulting in a 78 % increased MHB yield of 4.8 mol per mole of glucose. The strain $\Delta gapA/pEKEx2-Lbadh$ showed a 62 % decreased MHB production rate and an 87 % reduced glucose consumption rate, corresponding to a 193 % increase of the MHB yield of 7.9 mol per mole of glucose. As discussed below, the strongly reduced glucose uptake rate of the strain $\Delta gapA/pEKEx2-Lbadh$ is most likely a consequence of the fact that the strain does not form PEP.

By-product formation of wild-type and mutant strains

During biotransformation, by-product formation was nearly constant and specific rates were calculated (Table 5). The strain WT/ $pEKEx2-Lbadh$ showed an acetate formation rate (1.19 $mmol\ h^{-1}\ g_{cdw}^{-1}$) comparable to the glucose consumption rate (1.17 $mmol\ h^{-1}\ g_{cdw}^{-1}$). In addition, WT/ $pEKEx2-Lbadh$ formed succinate as by-product with a rate of 0.19 $mmol\ h^{-1}\ g_{cdw}^{-1}$. A low acetate production rate of 0.05 $mmol\ h^{-1}\ g_{cdw}^{-1}$ was shown by the strain $\Delta pfkA/pEKEx2-Lbadh$, which corresponds to only 8 % of the glucose uptake rate. Succinate was not formed by $\Delta pfkA/pEKEx2-Lbadh$. The strain $\Delta gapA/pEKEx2-Lbadh$ formed neither acetate nor succinate, but glycerol with a rate of 0.08 $mmol\ h^{-1}\ g_{cdw}^{-1}$, which corresponds to 53 % of the glucose consumption rate. As glyceraldehyde 3-phosphate cannot be catabolized to pyruvate in the $\Delta gapA$ mutant, reduction to glycerol presents an alternative pathway to oxidation in the cyclic PPP.

Discussion

For reductive whole-cell biotransformations requiring NADPH, attempts were made in this work to increase the NADPH yield per mole of glucose using *C. glutamicum* as host strain and the reduction of MAA to MHB as NADPH-requiring model reaction. Rerouting of glucose catabolism from glycolysis to the oxidative PPP was achieved by deletion of either the *pfkA* gene or the *gapA* gene.

C. glutamicum wild type carrying $pEKEx2-Lbadh$ showed a 31 % lower specific MHB production rate compared to *E. coli* carrying $pBtac-Lbadh$, even when compared to an *E. coli* biotransformation conducted at 30 °C (unpublished data). This difference might be due to a lower glucose uptake capacity or to a generally lower metabolic flux capacity of *C. glutamicum*. Overexpression of the genes involved in glucose uptake and catabolism via glycolysis

Table 4 Growth rates (μ) and biomass concentrations [cell dry weight (cdw) l^{-1}] in glucose minimal medium with 1 mM IPTG and 100 $\mu g\ ml^{-1}$ spectinomycin, and specific NAD^+ -dependent glyceraldehyde 3-phosphate dehydrogenase (GAPDH) activity in cell extracts of the indicated *C. glutamicum* strains after cultivation in LB medium with 1 mM IPTG and 100 $\mu g\ ml^{-1}$ spectinomycin

<i>C. glutamicum</i>	$\mu\ (h^{-1})$	$cdw\ (g\ l^{-1})^a$	GAPDH activity ($\mu mol\ min^{-1}\ mg^{-1}$)
WT/ $pEKEx3$	0.33±0.01	7.80±0.07	0.15±0.02
WT/ $pEKEx3-gapA^{Cgl}$	0.31±0.00	8.08±0.11	0.26±0.03
$\Delta gapA/pEKEx3$	0.00±0.01	0.00±0.00 ^b	0.00±0.00
$\Delta gapA/pEKEx3-gapA^{Cgl}$	0.27±0.01	7.99±0.30	0.13±0.02

^a Determination of cdw at maximal biomass

^b Determination of cdw after 24 h




Fig. 2 Kinetics of MHB production (open squares) and glucose consumption (filled squares) during biotransformation of MAA to MHB using resting cells ($3 \text{ g}_{\text{cdw}} \text{ l}^{-1}$) of the indicated *C. glutamicum* strains

or PPP could improve the rate of glucose catabolism, as shown recently for oxygen-deprived conditions (Yamamoto et al. 2012; Jojima et al. 2010). The MHB per glucose yield found for *C. glutamicum* WT/pEKEx2-*Lbadh* (2.7 mol/mol) was 10 % higher than the corresponding value determined for *E. coli* BL21(DE3)/pBtac-*Lbadh* (2.44 mol/mol) (Siedler et al. 2011), which might be due to slight differences in the partition of glucose 6-phosphate between glycolysis and the PPP.

Biotransformation studies with *E. coli* $\Delta pfkA$ and $\Delta pfkA \Delta pfkB$ mutants expressing *Lbadh* showed yields of 4.8 and $5.4 \text{ mol}_{\text{MHB}} \text{ mol}_{\text{glucose}}^{-1}$, respectively (Siedler et al. 2011). ^{13}C metabolic flux analysis demonstrated a negative net flux through phosphoglucose isomerase in the $\Delta pfkA$ mutant, in compliance with the proposed partial cyclization of the PPP (Siedler et al. 2012). The MHB yield per glucose of the *E. coli* strain $\Delta pfkA$ /pBtac-*Lbadh* was comparable to that of the *C. glutamicum* strain $\Delta pfkA$ /pEKEx2-*Lbadh* ($4.8 \text{ mol}_{\text{MHB}} \text{ mol}_{\text{glucose}}^{-1}$), indicating that a partial cyclization of the PPP occurred in the latter species, too. Furthermore, similarities were found when comparing by-product formation in *E. coli* and *C. glutamicum*. Less acetate and no succinate was produced in both $\Delta pfkA$ mutant strains compared to the reference strains within the experimental period, presumably as a consequence of a decreased carbon flux through the lower part of glycolysis and the TCA cycle in these mutants (Siedler et al. 2012).

C. glutamicum possesses two glyceraldehyde 3-phosphate dehydrogenases (GAPDH), but only GapA functions in the glycolytic direction (Omumasaba et al. 2004). Thus, a

deletion of the corresponding gene theoretically should result in a cyclization of the PPP. The fact that the MHB per glucose yield of the strain $\Delta gapA$ /pEKEx2-*Lbadh* (7.9 mol/mol) was higher compared to the strain $\Delta pfkA$ /pEKEx2-*Lbadh* and corresponded to 66 % of the maximal value of 12 mol NADPH per mole of glucose indicated a more extended cyclic operation of the PPP in the $\Delta gapA$ mutant compared to the $\Delta pfkA$ mutant. The maximal value for a complete oxidation of glucose in the PPP was not reached because 25 % of the glucose carbon was lost by reduction of glyceraldehyde 3-phosphate to glycerol. Taking this loss into account, only $9 \text{ mol}_{\text{MHB}} \text{ mol}_{\text{glucose}}^{-1}$ could be achieved maximally. The experimental yield of $7.9 \text{ mol}_{\text{MHB}} \text{ mol}_{\text{glucose}}^{-1}$ corresponds to 88 % of this value and is 46 % above the best yields reported so far (Chin and Cirino 2011; Siedler et al. 2011, 2012). Future yield optimization could be achieved by deletion of the gene encoding glycerol 3-phosphatase. Such a deletion was recently shown to prevent glycerol formation, which predominantly occurs in fructose-utilizing *C. glutamicum* strains (Lindner et al. 2012).

The strongly reduced biotransformation rate of the strain $\Delta gapA$ /pEKEx2-*Lbadh* was probably a consequence of the diminished capability for glucose uptake. In a $\Delta gapA$ mutant, no PEP should be formed during glucose catabolism and consequently, glucose uptake via the PTS should be impossible. PTS-independent glucose uptake has recently been described for *C. glutamicum*. It involves the inositol transporters IolT1 and IolT2 which also function as low-affinity glucose permeases (Lindner et al. 2011). Subsequent phosphorylation of glucose to glucose 6-phosphate is

The strongly reduced biotransformation rate of the strain $\Delta gapA$ /pEKEx2-*Lbadh* was probably a consequence of the diminished capability for glucose uptake. In a $\Delta gapA$ mutant, no PEP should be formed during glucose catabolism and consequently, glucose uptake via the PTS should be impossible. PTS-independent glucose uptake has recently been described for *C. glutamicum*. It involves the inositol transporters IolT1 and IolT2 which also function as low-affinity glucose permeases (Lindner et al. 2011). Subsequent phosphorylation of glucose to glucose 6-phosphate is

Table 5 Biotransformation parameters and by-product formation of *C. glutamicum* wild-type and deletion mutants carrying plasmid pEKEx2-*Lbadh*

<i>C. glutamicum</i> strain	Specific MHB production rate ($\text{mmol h}^{-1} \text{ g}_{\text{cdw}}^{-1}$)	Specific glucose consumption rate ($\text{mmol h}^{-1} \text{ g}_{\text{cdw}}^{-1}$)	Yield ($\text{mol}_{\text{MHB}} \text{ mol}_{\text{Glucose}}^{-1}$)	Specific acetate formation rate ($\text{mmol h}^{-1} \text{ g}_{\text{cdw}}^{-1}$)	Specific succinate formation rate ($\text{mmol h}^{-1} \text{ g}_{\text{cdw}}^{-1}$)	Specific glycerol formation rate ($\text{mmol h}^{-1} \text{ g}_{\text{cdw}}^{-1}$)
WT/pEKEx2- <i>Lbadh</i>	3.14 ± 0.13	1.17 ± 0.07	2.7 ± 0.1	1.19 ± 0.01	0.19 ± 0.01	0
$\Delta pfkA$ /pEKEx2- <i>Lbadh</i>	2.88 ± 0.08	0.60 ± 0.01	4.8 ± 0.2	0.05 ± 0.01	0	0
$\Delta gapA$ /pEKEx2- <i>Lbadh</i>	1.20 ± 0.04	0.15 ± 0.03	7.9 ± 0.9	0	0	0.08 ± 0.04

catalyzed either by an ATP-dependent glucokinase encoded by *glk* (Park et al. 2000) or by the polyphosphate- or ATP-dependent glucose kinase PpgK (Lindner et al. 2010). It can be assumed that glucose uptake during biotransformation with the $\Delta gapA$ mutant occurs via this alternative pathway, as the observed glucose consumption rate of $2.5 \text{ nmol min}^{-1} \text{ mg}_{\text{cdw}}^{-1}$ (Table 5) at glucose concentrations >10-fold above the apparent K_s values of IolT1 and IolT2 (2.8 and 1.9 mM, respectively) is in the range determined for PTS-independent glucose uptake at 1 mM glucose ($0.7 \text{ nmol min}^{-1} \text{ mg}_{\text{cdw}}^{-1}$) (Lindner et al. 2011). Overexpression of either *iolT1* or *iolT2* together with *ppgK* was shown to allow almost wild-type growth rates in a PTS-negative mutant (Lindner et al. 2011) and thus would probably also allow higher biotransformation rates of a $\Delta gapA$ mutant. Alternatively, expression of the glucose facilitator gene *glf* from *Zymomonas mobilis* could help to increase glucose uptake (Weisser et al. 1995; Parker et al. 1995).

Overall, we could demonstrate the potential of *C. glutamicum* for NADPH-dependent reductive whole-cell biotransformation and show that deletion of either *pfkA* or *gapA* is beneficial to improve the NADPH per glucose yield, presumably by cyclization of the PPP.

Acknowledgments This work was supported by the Ministry of Innovation, Science, Research and Technology of North Rhine-Westphalia (BioNRW, Technology Platform Biocatalysis, RedoxCell support code—W0805wb001b).

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

- Abe S, Takayama K, Kinoshita S (1967) Taxonomical studies on glutamic acid producing bacteria. *J Gen Appl Microbiol* 13 (3):279–301
- Akinterinwa O, Cirino PC (2011) Anaerobic obligatory xylitol production in *Escherichia coli* strains devoid of native fermentation pathways. *Appl Environ Microbiol* 77(2):706–709
- Babul J (1978) Phosphofructokinases from *Escherichia coli*. Purification and characterization of the nonallosteric isozyme. *J Biol Chem* 253 (12):4350–4355
- Bäumchen C, Bringer-Meyer S (2007) Expression of *glf*_{Zm} increases D-mannitol formation in whole cell biotransformation with resting cells of *Corynebacterium glutamicum*. *Appl Microbiol Biotechnol* 76(3):545–552
- Becker J, Wittmann C (2011) Bio-based production of chemicals, materials and fuels—*Corynebacterium glutamicum* as versatile cell factory. *Curr Opin Biotechnol*. doi:10.1016/j.copbio.2011.11.012
- Blombach B, Riester T, Wieschalka S, Ziert C, Youn JW, Wendisch VF, Eikmanns BJ (2011) *Corynebacterium glutamicum* tailored for efficient isobutanol production. *Appl Environ Microbiol* 77 (10):3300–3310
- Chemler JA, Fowler ZL, McHugh KP, Koffas MA (2010) Improving NADPH availability for natural product biosynthesis in *Escherichia coli* by metabolic engineering. *Metab Eng* 12(2):96–104
- Chin JW, Cirino PC (2011) Improved NADPH supply for xylitol production by engineered *Escherichia coli* with glycolytic mutations. *Biotechnol Prog* 27(2):333–341
- Doo EH, Lee WH, Seo HS, Park JB (2009) Productivity of cyclohexanone oxidation of the recombinant *Corynebacterium glutamicum* expressing *chnB* of *Acinetobacter calcoaceticus*. *J Biotechnol* 142(2):164–169
- Eggeling L, Bott M (eds) (2005) Handbook of *Corynebacterium glutamicum*. CRC, Boca Raton, pp 535–568
- Eguchi T, Kuge Y, Inoue K, Yoshikawa N, Mochida K, Uwajima T (1992) NADPH regeneration by glucose dehydrogenase from *Gluconobacter scleroides* for l-leucovorin synthesis. *Biosci Biotechnol Biochem* 56(5):701–703
- Eikmanns BJ, Kleinertz E, Liebl W, Sahm H (1991) A family of *Corynebacterium glutamicum*/*Escherichia coli* shuttle vectors for cloning, controlled gene expression, and promoter probing. *Gene* 102(1):93–98
- Eikmanns BJ, Rittmann D, Sahm H (1995) Cloning, sequence analysis, expression, and inactivation of the *Corynebacterium glutamicum* *icd* gene encoding isocitrate dehydrogenase and biochemical characterization of the enzyme. *J Bacteriol* 177(3):774–782
- Ernst M, Kaup B, Müller M, Bringer-Meyer S, Sahm H (2005) Enantioselective reduction of carbonyl compounds by whole-cell biotransformation, combining a formate dehydrogenase and a (R)-specific alcohol dehydrogenase. *Appl Microbiol Biotechnol* 66 (6):629–634
- Fasan R, Crook NC, Peters MW, Meinhold P, Buelter T, Landwehr M, Cirino PC, Arnold FH (2011) Improved product-per-glucose yields in P450-dependent propane biotransformations using engineered *Escherichia coli*. *Biotechnol Bioeng* 108(3):500–510
- Hanahan D (1983) Studies on transformation of *Escherichia coli* with plasmids. *J Mol Biol* 166(4):557–580
- Hanahan D, Jessee J, Bloom FR (1991) Plasmid transformation of *Escherichia coli* and other bacteria. *Methods Enzymol* 204:63–113
- Hermann T (2003) Industrial production of amino acids by coryneform bacteria. *J Biotechnol* 104(1–3):155–172
- Inui M, Kawaguchi H, Murakami S, Vertes AA, Yukawa H (2004) Metabolic engineering of *Corynebacterium glutamicum* for fuel ethanol production under oxygen-deprivation conditions. *J Mol Microbiol Biotechnol* 8(4):243–254
- Ishige T, Honda K, Shimizu S (2005) Whole organism biocatalysis. *Curr Opin Chem Biol* 9(2):174–180
- Jojima T, Fujii M, Mori E, Inui M, Yukawa H (2010) Engineering of sugar metabolism of *Corynebacterium glutamicum* for production of amino acid L-alanine under oxygen deprivation. *Appl Microbiol Biotechnol* 87(1):159–165
- Kabus A, Niebisch A, Bott M (2007) Role of cytochrome *bd* oxidase from *Corynebacterium glutamicum* in growth and lysine production. *Appl Environ Microbiol* 73(3):861–868
- Kaup B, Bringer-Meyer S, Sahm H (2004) Metabolic engineering of *Escherichia coli*: construction of an efficient biocatalyst for D-mannitol formation in a whole-cell biotransformation. *Appl Microbiol Biotechnol* 64(3):333–339
- Kaup B, Bringer-Meyer S, Sahm H (2005) D-Mannitol formation from D-glucose in a whole-cell biotransformation with recombinant *Escherichia coli*. *Appl Microbiol Biotechnol* 69(4):397–403
- Kimura E (2003) Metabolic engineering of glutamate production. *Adv Biochem Eng Biotechnol* 79:37–57
- Kruger NJ, von Schaewen A (2003) The oxidative pentose phosphate pathway: structure and organisation. *Curr Opin Plant Biol* 6 (3):236–246
- Lindner SN, Knebel S, Pallerla SR, Schoberth SM, Wendisch VF (2010) Cg2091 encodes a polyphosphate/ATP-dependent glucokinase of *Corynebacterium glutamicum*. *Appl Microbiol Biotechnol* 87(2):703–713

- Lindner SN, Seibold GM, Henrich A, Krämer R, Wendisch VF (2011) Phosphotransferase system-independent glucose utilization in *Corynebacterium glutamicum* by inositol permeases and glucokinases. *Appl Environ Microbiol* 77(11):3571–3581
- Lindner SN, Meiswinkel TM, Panhorst M, Youn JW, Wiefel L, Wendisch VF (2012) Glycerol-3-phosphatase of *Corynebacterium glutamicum*. *J Biotechnol*. doi:10.1016/j.jbiotec.2012.02.003
- Litsanov B, Brocker M, Bott M (2012a) Towards homosuccinate fermentation: metabolic engineering of *Corynebacterium glutamicum* for anaerobic succinate production from glucose and formate. *Appl Environ Microbiol*. doi:10.1128/AEM.07790-11
- Litsanov B, Kabus A, Brocker M, Bott M (2012b) Efficient aerobic succinate production from glucose in minimal medium with *Corynebacterium glutamicum*. *Microb Biotechnol* 5(1):116–128
- Meissner D, Vollstedt A, van Dijl JM, Freudl R (2007) Comparative analysis of twin-arginine (Tat)-dependent protein secretion of a heterologous model protein (GFP) in three different Gram-positive bacteria. *Appl Microbiol Biotechnol* 76(3):633–642
- Miller JH (ed) (1972) Experiments in molecular genetics. Cold Spring Harbor Laboratory, New York
- Mimitsuka T, Sawai H, Hatsu M, Yamada K (2007) Metabolic engineering of *Corynebacterium glutamicum* for cadaverine fermentation. *Biosci Biotechnol Biochem* 71(9):2130–2135
- Niebisch A, Bott M (2001) Molecular analysis of the cytochrome *bc₁-aa₃* branch of the *Corynebacterium glutamicum* respiratory chain containing an unusual diheme cytochrome *c₁*. *Arch Microbiol* 175(4):282–294
- Okino S, Noburyu R, Suda M, Jojima T, Inui M, Yukawa H (2008) An efficient succinic acid production process in a metabolically engineered *Corynebacterium glutamicum* strain. *Appl Microbiol Biotechnol* 81(3):459–464
- Omumasaba CA, Okai N, Inui M, Yukawa H (2004) *Corynebacterium glutamicum* glyceraldehyde-3-phosphate dehydrogenase isoforms with opposite, ATP-dependent regulation. *J Mol Microbiol Biotechnol* 8(2):91–103
- Panke S, Wubbolts M (2005) Advances in biocatalytic synthesis of pharmaceutical intermediates. *Curr Opin Chem Biol* 9(2):188–194
- Park SY, Kim HK, Yoo SK, Oh TK, Lee JK (2000) Characterization of *glk*, a gene coding for glucose kinase of *Corynebacterium glutamicum*. *FEMS Microbiol Lett* 188(2):209–215
- Parker C, Barnell WO, Snoep JL, Ingram LO, Conway T (1995) Characterization of the *Zymomonas mobilis* glucose facilitator gene product (*glf*) in recombinant *Escherichia coli*: examination of transport mechanism, kinetics and the role of glucokinase in glucose transport. *Mol Microbiol* 15(5):795–802
- Pfefferle W, Möckel B, Bathe B, Marx A (2003) Biotechnological manufacture of lysine. *Adv Biochem Eng Biotechnol* 79:59–112
- Rittmann D, Schaffer S, Wendisch VF, Sahm H (2003) Fructose-1, 6-bisphosphatase from *Corynebacterium glutamicum*: expression and deletion of the *fbp* gene and biochemical characterization of the enzyme. *Arch Microbiol* 180(4):285–292
- Sambrook J, Russell DW (eds) (2001) Molecular cloning: a laboratory manual. Cold Spring Harbor Laboratory, Cold Spring Harbor
- Schäfer A, Tauch A, Jäger W, Kalinowski J, Thierbach G, Pühler A (1994) Small mobilizable multi-purpose cloning vectors derived from the *Escherichia coli* plasmids pK18 and pK19: selection of defined deletions in the chromosome of *Corynebacterium glutamicum*. *Gene* 145(1):69–73
- Schneider J, Wendisch VF (2010) Putrescine production by engineered *Corynebacterium glutamicum*. *Appl Microbiol Biotechnol* 88(4):859–868
- Siedler S, Bringer S, Bott M (2011) Increased NADPH availability in *Escherichia coli*: improvement of the product per glucose ratio in reductive whole-cell biotransformation. *Appl Microbiol Biotechnol* 92(5):929–937
- Siedler S, Bringer S, Blank LM, Bott M (2012) Engineering yield and rate of reductive biotransformation in *Escherichia coli* by partial cyclization of the pentose phosphate pathway and PTS-independent glucose transport. *Appl Microbiol Biotechnol* 93(4):1459–1467
- Smith KM, Cho KM, Liao JC (2010) Engineering *Corynebacterium glutamicum* for isobutanol production. *Appl Microbiol Biotechnol* 87(3):1045–1055
- Stäbler N, Oikawa T, Bott M, Eggeling L (2011) *Corynebacterium glutamicum* as a host for synthesis and export of D-amino acids. *J Bacteriol* 193(7):1702–1709
- Stansen C, Uy D, Delaunay S, Eggeling L, Goergen JL, Wendisch VF (2005) Characterization of a *Corynebacterium glutamicum* lactate utilization operon induced during temperature-triggered glutamate production. *Appl Environ Microbiol* 71(10):5920–5928
- Tan I (2006) Applications of whole cell biotransformations for the production of chiral alcohols. Rheinische Friedrich-Wilhelms University of Bonn, Bonn, Dissertation
- Weisser P, Krämer R, Sahm H, Sprenger GA (1995) Functional expression of the glucose transporter of *Zymomonas mobilis* leads to restoration of glucose and fructose uptake in *Escherichia coli* mutants and provides evidence for its facilitator action. *J Bacteriol* 177(11):3351–3354
- Wendisch VF, Bott M, Eikmanns BJ (2006) Metabolic engineering of *Escherichia coli* and *Corynebacterium glutamicum* for biotechnological production of organic acids and amino acids. *Curr Opin Microbiol* 9(3):268–274
- Yamamoto S, Gunji W, Suzuki H, Toda H, Suda M, Jojima T, Inui M, Yukawa H (2012) Overexpression of glycolytic genes enhances *Corynebacterium glutamicum* glucose metabolism and alanine production under oxygen-deprived conditions. *Appl Environ Microbiol*. doi:10.1128/AEM.07998-11
- Yokota A, Lindley ND (2005) Central metabolism: sugar uptake and conversion. In: Eggeling L, Bott M (eds) *Handbook of Corynebacterium glutamicum*. CRC, Boca Raton, pp 215–240
- Yun JY, Lee JE, Yang KM, Cho S, Kim A, Kwon YE, Park JB (2012) Ethambutol-mediated cell wall modification in recombinant *Corynebacterium glutamicum* increases the biotransformation rates of cyclohexanone derivatives. *Bioprocess Biosyst Eng* 35(1–2):211–216